



Track your LEQEMBI IQLIK™ injections



You can use this tracker to help stay on top of your LEQEMBI IQLIK (pronounced "i-click") injection schedule.



LEQEMBI Prescriber	
Name	
Phone	
Phone	
Specialty Pharmacy	
Name	
Phone	





My LEQEMBI at-home injection tracker



When injecting, refer to the Instructions for Use. You can find it in the LEQEMBI IQLIK box or at **LEQEMBI.com/injection**.



You can also visit **LEQEMBI.com/HowToInject** for an overview of the injection process.

Write down the date and time, and circle the location of your weekly injection Example Date TUES 7/15 Time 2:30 pm Notes Circle the location on your body where you injected.



WHAT IS LEQEMBI?

LEQEMBI is a prescription medicine used to treat people with early Alzheimer's disease, which includes mild cognitive impairment (MCI) or mild dementia stage of disease.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about LEQEMBI?

LEQEMBI can cause serious side effects, including: ARIA (Amyloid-Related Imaging Abnormalities). ARIA is a side effect that does not usually cause any symptoms, but serious symptoms can occur. ARIA can be fatal. ARIA commonly shows up as temporary swelling in areas of the brain that usually goes away over time. Small spots of bleeding in or on the surface of the brain can occur. Less often, larger areas of bleeding in the brain can occur. Most people with ARIA don't have any symptoms. However, some people may notice: headache, confusion that gets worse, dizziness, vision changes, nausea, difficulty walking, seizures, difficulty speaking, or muscle weakness.

- Some people have a gene called ApoE4 that may increase the risk of ARIA. Talk to your healthcare provider about testing to see if you have this gene
- You may be at a higher risk of developing bleeding in the brain if you take medicines to reduce blood clots from forming (antithrombotic medicines) while receiving LEQEMBI. Talk to your healthcare provider to see if any of the medicines you're taking increase this risk
- Your healthcare provider will check for ARIA with MRI (magnetic resonance imaging) scans before you start LEQEMBI and during treatment
- You should carry information that says you are receiving LEQEMBI, which can cause ARIA, and that ARIA symptoms can look like stroke symptoms

Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the symptoms listed above.

Serious allergic reactions:

Do not receive LEQEMBI if you have serious allergic reactions to LEQEMBI, LEQEMBI IQLIK, or any of the ingredients.

• Tell your healthcare provider if you notice any symptoms during or after a LEQEMBI infusion, including: swelling of the face, lips, mouth, or tongue; itchy bumps on the skin, also known as hives; or difficulty breathing

Infusion-related reactions:

- Infusion-related reactions can occur during or after completion with LEQEMBI injection into a vein (intravenously), which can be serious. Tell your healthcare provider right away if you notice any of these symptoms: fever; flu-like symptoms (chills, body aches, feeling shaky, joint pain); nausea and/or vomiting; dizziness or lightheadedness; fast or slow heart rate, or feeling like your chest is pounding; difficulty breathing or shortness of breath; and changes in blood pressure
- If you have an infusion-related reaction, your healthcare provider may give you medicines before your next infusion to lower the chance of having a reaction

Injection-related reactions:

• Injection-related reactions may occur with LEQEMBI injection under the skin (subcutaneous injection with LEQEMBI IQLIK). Tell your healthcare provider if you notice any of these symptoms during or after an injection: redness, swelling, heat, pain, itching, rash, bruising, and blood collection under the skin at the injection site; headache, fatigue, or fever may also be observed after an injection

The most common side effects of LEQEMBI include infusion-related reactions, ARIA, and headaches.

These are not all the possible side effects of LEQEMBI. Call your doctor for more information and medical advice about side effects. You may report side effects to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

Before receiving LEQEMBI, tell your healthcare provider about:


- **All your medical conditions**, including if you are pregnant, breastfeeding, or plan to become pregnant or breastfeed. It is not known if LEQEMBI could harm your unborn or breastfeeding baby
- All the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take medicines to reduce blood clots from forming (antithrombotic medicines, including aspirin)

LEQEMBI (lecanemab-irmb) is available as:

- Intravenous infusion: 100 mg/mL
- Subcutaneous injection: 200 mg/mL



Where to inject



Date	
Time	
Notes	

Time Notes	
Date Time	
Notes	

Date	
Notes	

Date	
Time	
Notes	


Date	
Time	
Notes	

Date	
Time	
Notes	

Date Time	
Notes	
Date	

Time

Notes



Date	
Time	
Notes	

Date	
Time	
Notes	

Date	
Time	
Notes	

Date	
Time	
Notes	

Date Time	
Notes	
Date Time	
Notes	

Date	
Time	
Notes	

Date Time	
Notes	

Date	
Time	
Notes	

Date	
Time	
Notes	

Date Time	
Notes	
Date	
Time	
Notes	

Notes	

Date	
Time	
Notes	

Date Time	
Notes	

Date	
Time	
Notes	

Time Notes	
Date Time	
Notes	



Want to track injections on your mobile phone or tablet?

The LEQEMBI Companion™ app can help.

Visit <u>LEQEMBI.com/CompanionAppSignUp</u> to learn how to get started

You do not need to be enrolled in the LEQEMBI Companion $^{\!\scriptscriptstyle\mathsf{TM}}$ program to download and use the app.

Please see Important Safety Information on <u>page 4</u>. Click underlined links for <u>Medication Guide with Instructions for Use</u> and <u>full Prescribing Information</u>, including Boxed WARNING.

