

STARTING TREATMENT

with LEQEMBI® IV INFUSIONS

Getting started and
staying the course

NINA

LEQEMBI PATIENT SINCE 2023

People shown were compensated for their time, and information is accurate as of August 2026.

WHAT IS LEQEMBI?

LEQEMBI is a prescription medicine used to treat people with early Alzheimer's disease, which includes mild cognitive impairment (MCI) or mild dementia stage of disease.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about LEQEMBI?

LEQEMBI can cause serious side effects, including: ARIA (Amyloid-Related Imaging Abnormalities). ARIA is a side effect that does not usually cause any symptoms, but serious symptoms can occur. ARIA can be fatal.

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.

 **LEQEMBI®**
(lecanemab-irmb)

NAVIGATING YOUR TREATMENT JOURNEY

Even though starting treatment can feel overwhelming, this guide is here to help you.

It provides useful information at every stage: while you're starting treatment, when you need support, and to help you stay on track.

ABOUT LEQEMBI IV TREATMENT

An overview for starting your treatment



Given as intravenous (IV) infusions

This means a needle is placed in a vein in your arm to give the medicine. Each infusion takes about 1 hour.



At a healthcare provider's office or infusion center

Talk to your healthcare provider and insurance provider about where to get your infusions.



Receive treatment twice monthly (once every 2 weeks)

If you miss an infusion of LEQEMBI, reach out to your healthcare provider. You should receive your next dose as soon as possible.



LEQEMBI is the only anti-amyloid treatment with maintenance dosing options

After 18 months of infusions every 2 weeks, you and your healthcare provider will decide if you can continue taking LEQEMBI or switch to a less frequent dose to keep slowing the progression of early Alzheimer's disease.

Prefer to treat at home or on the go?*

Ask your healthcare provider if self-injections with the LEQEMBI IQLIK® autoinjector may be an option for you.

*For information on storing and using LEQEMBI IQLIK, please see Instructions for Use on [LEQEMBI.com/instructionsforuse](https://www.leqembi.com/instructionsforuse). Once the LEQEMBI IQLIK autoinjector has been stored at room temperature, do not return it to the refrigerator.

IMPORTANT SAFETY INFORMATION (continued)

- ARIA commonly shows up as temporary swelling in areas of the brain that usually goes away over time
- Small spots of bleeding in or on the surface of the brain can occur
- Less often, larger areas of bleeding in the brain can occur
- Most people with ARIA don't have any symptoms. However, some people may notice: headache, confusion, dizziness, vision changes, nausea, difficulty walking, seizures, difficulty speaking, or muscle weakness



Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

ABOUT LEQEMBI IV TREATMENT

Helpful tips and information to prepare for your infusion



Keep your phone close by

Expect a call from your infusion center before your appointment. Ask about anything you need to bring, such as paperwork and medication lists.



Wear short sleeves

It'll make it easier to get your infusion at your appointment, but bring a blanket in case you get chilly.



Take along something to play, read, or watch

It will help make your appointment more enjoyable.



Bring snacks and a bottle of water

Drink plenty beforehand so you're hydrated.



Write down your questions

Bring a list of your concerns to discuss with your healthcare provider.

It can be helpful to have a family member with you during treatment, so ask your infusion center if that's allowed.

DOYLE

LEQEMBI PATIENT SINCE 2023

People shown were compensated for their time, and information is accurate as of August 2026.

IMPORTANT SAFETY INFORMATION (continued)

- Some people have a gene called ApoE4 that may increase the risk of ARIA. Talk to your healthcare provider about testing to see if you have this gene
- You may be at a higher risk of developing bleeding in the brain if you take medicines to reduce blood clots from forming (antithrombotic medicines) while receiving LEQEMBI. Talk to your healthcare provider to see if any of the medicines you're taking increase this risk

Monitoring for potential side effects

It's important to know that all anti-amyloid treatments for Alzheimer's disease require MRI (magnetic resonance imaging) monitoring. **LEQEMBI can cause serious side effects, including one called ARIA (Amyloid-Related Imaging Abnormalities).** ARIA commonly shows up as temporary swelling or bleeding in the brain, and can only be seen with MRI monitoring.

Most of the time, ARIA does not cause any symptoms, though serious symptoms can occur and can be fatal. **That's why you will receive an MRI before treatment and 4 MRIs after you start treatment—to help your healthcare provider spot any changes early.**

Your exact MRI schedule will be determined by your healthcare provider and may include additional MRIs.

Symptoms of ARIA

Most people with ARIA don't have any symptoms. However, some people may notice:



headache



vision changes



seizures



confusion



nausea



difficulty speaking



dizziness



difficulty walking



muscle weakness



Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

THE INFUSION PROCESS

A LEQEMBI IV infusion often includes these steps

1

A member of the infusion care team will check your vital signs.

You may also get weighed and have blood drawn.

2

Your care team will prepare your medicine, set up your IV line, and start your infusion.

Pay attention to how you feel and keep your care team informed.

3

After your infusion is finished, your care team will remove the IV.

They'll tell you when you can leave and how to schedule your next infusion.

If you have an infusion-related reaction

Your healthcare provider may:

- Adjust the flow or stop the infusion
- Give you medicines before your future LEQEMBI infusions to decrease your chance of having an infusion-related reaction. These medicines may include antihistamines, anti-inflammatory medicines, or steroids
- Stop your infusion entirely. Serious infusion reactions can also occur after completion of a LEQEMBI infusion

Tell your healthcare provider right away if you experience these symptoms during or after a LEQEMBI infusion.

IMPORTANT SAFETY INFORMATION (continued)

- Your healthcare provider will check for ARIA with MRI (magnetic resonance imaging) scans before you start LEQEMBI and during treatment
- You should carry information that says you are receiving LEQEMBI, which can cause ARIA, and that ARIA symptoms can look like stroke symptoms

Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the symptoms listed on page 3.



For care partners

To help make these appointments more enjoyable, consider games or activities you can do together.

KAMALA

LEQEMBI PATIENT SINCE 2024

People shown were compensated for their time, and information is accurate as of August 2026.

Serious allergic reactions:

Do not receive LEQEMBI if you have serious allergic reactions to LEQEMBI, LEQEMBI IQLIK, or any of the ingredients.

- Tell your healthcare provider and get medical help right away if you notice any symptoms during or after treatment with LEQEMBI, including: swelling of the face, lips, mouth, or tongue; itchy bumps on the skin, also known as hives; or difficulty breathing

 **LEQEMBI**[®]
(lecanemab-irmb)

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

NAVIGATING TREATMENT

Practical tips for staying on track

In people with Alzheimer's disease, harmful amyloid brain plaque continuously builds up. LEQEMBI keeps working throughout treatment to remove it.

That's why it's important to stick with your treatment plan.



Schedule upcoming infusions in advance

Add the appointments, including follow-up MRIs, to your calendar.



Track how you're feeling

Take notes between your infusions and keep your healthcare provider informed of any updates.



Missed an infusion?

Contact your healthcare provider so they can help you get back on track.

LEQEMBI can cause serious side effects

These include ARIA, serious allergic reactions, and infusion- or injection-related reactions.

The most common side effects of LEQEMBI

These include infusion-related reactions with LEQEMBI injected into a vein, ARIA, and headache.

These are not all the possible side effects of LEQEMBI and LEQEMBI IQLIK. For more information, ask your healthcare provider or pharmacist.

IMPORTANT SAFETY INFORMATION (continued)

Infusion-related reactions:

- Infusion-related reactions are a common side effect that can occur during or after completion with LEQEMBI injection into a vein (intravenously), which can be serious. Tell your healthcare provider right away if you notice any of these symptoms: fever; flu-like symptoms (chills, body aches, feeling shaky, joint pain); nausea and/or vomiting; dizziness or lightheadedness; fast or slow heart rate, or feeling like your chest is pounding; difficulty breathing or shortness of breath; or changes in blood pressure
- If you have an infusion-related reaction, your healthcare provider may give you medicines before your next infusion to lower the chance of having a reaction

Ask about LEQEMBI IQLIK

Talk to your healthcare provider to find out if you can start treating at home or on the go.*



LEONARD

LEQEMBI PATIENT SINCE 2023

Patient shown was compensated for their time, and information is accurate as of August 2026.



If you and your healthcare provider decide you can continue receiving LEQEMBI after 18 months, treatment will be a once-monthly (once every 4 weeks) infusion or once-weekly self-injections with LEQEMBI IQLIK.

*For information on storing and using LEQEMBI IQLIK, please see Instructions for Use on [LEQEMBI.com/instructionsforuse](https://www.leqembi.com/instructionsforuse). Once the LEQEMBI IQLIK autoinjector has been stored at room temperature, do not return it to the refrigerator.

Injection-related reactions:

- Injection-related reactions may occur with LEQEMBI injection under the skin (subcutaneous injection with LEQEMBI IQLIK). Tell your healthcare provider if you notice any of these symptoms during or after an injection: redness, swelling, heat, pain, itching, rash, bump, bruising, and blood collection under the skin at the injection site; headache, chills, fatigue, or fever may also be observed after an injection



Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING.**

HELPFUL RESOURCES



Navigate treatment with support by your side

Help is here throughout every step of your treatment, but **your healthcare provider is your go-to resource** for any urgent concerns.

The LEQEMBI Companion® program can help you:



Get information about accessing treatment



Understand your insurance coverage and potential out-of-pocket costs



Identify financial support programs, including the LEQEMBI Copay Assistance Program, if you are eligible for them

Visit LEQEMBI.com/PatientSupport to learn more

Get your treatment wallet card

The wallet card informs healthcare providers or emergency workers that you or your loved one is taking an anti-amyloid treatment. Access it in 2 ways:




Scan the QR code with your smartphone to add the card directly to your phone. It will appear in your digital wallet. Or print your card by visiting LEQEMBI.com/wallet.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of LEQEMBI include infusion-related reactions with LEQEMBI injected into a vein, ARIA, and headache.

Stay on track with the LEQEMBI Companion app

Your digital partner designed to support you at every step of your treatment. The app makes it easy to log your doses, set reminders, stay on top of your schedule, and more. Talk to your healthcare provider if you have any questions about your treatment plan.



Download the app



These steps will direct you to the Medisafe app. From there, you will select LEQEMBI as your treatment.

- 1 Scan the QR code, then tap the **pop-up banner**
- 2 Tap **"Get"** or **"Install"** to download Medisafe
- 3 Follow the prompts and tap **"Get Started"**
- 4 If prompted, enter the **verification code RW8Q**



Scan the QR code to download the app or visit LEQEMBI.com/CompanionAppSignUp to learn more about what the app has to offer.

Access tracker pages

If you'd prefer physical pages on which you can log your treatments and appointments, you can find them on pages 12-14 of this guide, or print out additional copies at LEQEMBI.com/InfusionTracker.



Please see Important Safety Information throughout. Click underlined links for **Medication Guide with Instructions for Use** and **full Prescribing Information**, including **Boxed WARNING**.

TRACKING INFUSIONS

The following pages are designed to help you track your **treatment dates** and **key information** regarding your LEQEMBI IV infusions.

First, fill out the information below regarding your infusion center.

Infusion center information

Address

Phone

Parking details

Example

Infusion #

5

Date Tuesday, August 15

Time 2:30pm

Reminders Arrive 15 min early.
Talk to Nurse Sarah about how to plan
my infusions with my vacation coming up.

Notes for care team

Keep track of
your infusions.

Note the date and time
of your treatments.

Write down any
questions you have.

Note how you're feeling
in between treatments
to share with your
healthcare provider.

MY INFUSIONS

Infusion #

Date

Time

Reminders

Notes for care team

Infusion #

Date

Time

Reminders

Notes for care team

Infusion #

Date

Time

Reminders

Notes for care team

Infusion #

Date

Time

Reminders

Notes for care team

IMPORTANT SAFETY INFORMATION (continued)

These are not all the possible side effects of LEQEMBI. Call your doctor for more information and medical advice about side effects. You may report side effects to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.



MY INFUSIONS

Infusion #	
Date	
Time	
Reminders	
Notes for care team	

Infusion #	
Date	
Time	
Reminders	
Notes for care team	

Infusion #	
Date	
Time	
Reminders	
Notes for care team	

Infusion #	
Date	
Time	
Reminders	
Notes for care team	

Download more
infusion log pages
by visiting
LEQEMBI.com/InfusionTracker

MY INFUSIONS	
Infusion #	
Date	
Time	
Reminders	
Notes for care team	



Prefer to track treatment on your phone?
Scan the QR code to download the app or
visit LEQEMBI.com/CompanionAppSignUp



Please see Important Safety Information throughout. Click underlined links
for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#),
including **Boxed WARNING**.

“

You just have to
**REALIZE
WHAT'S POSSIBLE
AND ACT.**

DOYLE

LEQEMBI PATIENT SINCE 2023

People shown were compensated for their time, and information is accurate as of August 2026. Quotes reflect personal experiences, which may vary.



Scan the QR code to read frequently asked questions about LEQEMBI. They can help you have a more informed discussion with your healthcare provider.

IMPORTANT SAFETY INFORMATION (continued)

Before receiving LEQEMBI, tell your healthcare provider about:

- **All your medical conditions**, including if you are pregnant, breastfeeding, or plan to become pregnant or breastfeed. It is not known if LEQEMBI could harm your unborn or breastfeeding baby
- **All the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take medicines to reduce blood clots from forming (antithrombotic medicines, including aspirin)

LEQEMBI (lecanemab-irmb) is available as:

- Intravenous infusion: 100 mg/mL
- Subcutaneous injection: 200 mg/mL



Please see Important Safety Information throughout. Click underlined links for Medication Guide with Instructions for Use and full Prescribing Information, including **Boxed WARNING.**



© 2026 Eisai Inc. and Biogen.
LEQE-US5499 September 2026

All trademarks are the property of their respective owners.

LEQEMBI®, LEQEMBI IQLIK®, and LEQEMBI Companion® are registered trademarks of Eisai R&D Management Co., Ltd.

