



For weekly maintenance treatment after 18 months

WHAT TO EXPECT WITH LEQEMBI® IQLIK™

(pronounced “i-click”)



IMPORTANT SAFETY INFORMATION

What is the most important information I should know about LEQEMBI?

LEQEMBI can cause serious side effects, including: ARIA (Amyloid-Related Imaging Abnormalities). ARIA is a side effect that does not usually cause any symptoms, but serious symptoms can occur. ARIA can be fatal. ARIA commonly shows up as temporary swelling in areas of the brain that usually goes away over time. Small spots of bleeding in or on the surface of the brain can occur. Less often, larger areas of bleeding in the brain can occur. Most people with ARIA don't have any symptoms. However, some people may notice: headache, confusion that gets worse, dizziness, vision changes, nausea, difficulty walking, seizures, difficulty speaking, or muscle weakness.

Please see Important Safety Information throughout. Click [underlined links](#) for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection



In this guide

Learn about maintenance treatment options, information about LEQEMBI IQLIK, and support available to you at every step.



Not actual size.

Actor portrayal.

WHAT IS LEQEMBI?

LEQEMBI is a prescription medicine used to treat people with early Alzheimer's disease, which includes mild cognitive impairment (MCI) or mild dementia stage of disease.

IMPORTANT SAFETY INFORMATION (CONTINUED)

What is the most important information I should know about LEQEMBI?

- Some people have a gene called ApoE4 that may increase the risk of ARIA. Talk to your healthcare provider about testing to see if you have this gene

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection



What is maintenance treatment?

Only LEQEMBI has FDA-approved maintenance dosing options to keep slowing progression

After receiving LEQEMBI infusions for 18 months, you may have the option of starting maintenance treatment with less frequent infusions or switching to injections.



Switch to
**ONCE-WEEKLY,
AT-HOME INJECTIONS**
(once every 7 days)

OR



Switch to
**ONCE-MONTHLY
INFUSIONS**
(once every 4 weeks)

Alternatively, you may continue on your twice-monthly (once every 2 weeks) infusion schedule if you and your healthcare provider determine that's the best plan for you.



It's important to stick to your treatment schedule. Your injections or infusions keep LEQEMBI working in your body and fighting Alzheimer's disease.

IMPORTANT SAFETY INFORMATION (CONTINUED)

What is the most important information I should know about LEQEMBI?


- You may be at a higher risk of developing bleeding in the brain if you take medicines to reduce blood clots from forming (antithrombotic medicines) while receiving LEQEMBI. Talk to your healthcare provider to see if any of the medicines you're taking increase this risk
- Your healthcare provider will check for ARIA with MRI (magnetic resonance imaging) scans before you start LEQEMBI and during treatment
- You should carry information that says you are receiving LEQEMBI, which can cause ARIA, and that ARIA symptoms can look like stroke symptoms

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection

What is LEQEMBI IQLIK?

LEQEMBI IQLIK is a single-dose, pen-like autoinjector that you or a care partner use to inject LEQEMBI under your skin (a subcutaneous injection)



Pen-like autoinjector

The needle tip is capped before injection. You inject by simply pushing the autoinjector against your skin.

No need to measure

The single-use autoinjector is prefilled to ensure you have the right dose.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the symptoms listed on the front cover.

Serious allergic reactions:

Do not receive LEQEMBI if you have serious allergic reactions to LEQEMBI, LEQEMBI IQLIK, or any of the ingredients.

- Tell your healthcare provider if you notice any symptoms during or after a LEQEMBI infusion, including:
 - swelling of the face, lips, mouth, or tongue
 - itchy bumps on the skin, also known as hives
 - difficulty breathing

Please see Important Safety Information throughout. Click [underlined links](#) for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection

Why choose LEQEMBI IQLIK?



Takes **less time** to complete than infusions



Can be given in the comfort and **privacy of your own home**



You can travel and still stay on schedule

When will I start LEQEMBI IQLIK?

After 18 months on LEQEMBI, your first at-home injection for maintenance treatment should be 2 weeks from your last infusion. If you've already started maintenance treatment with infusions, it should be 1 week from your last infusion. Your healthcare provider can help answer any questions you may have about starting LEQEMBI IQLIK.



Once your healthcare provider prescribes LEQEMBI IQLIK, your specialty pharmacy will verify your insurance coverage. You should **expect a call from the specialty pharmacy** (it may show as an unknown number) to confirm **where and when your medicine should be delivered**. Once you receive your LEQEMBI IQLIK, be sure to store it in the refrigerator until you are ready to inject.



Actor portrayal.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Infusion-related reactions:

- Infusion-related reactions can occur during or after completion with LEQEMBI injection into a vein (intravenously), which can be serious. Tell your healthcare provider right away if you notice any of these symptoms:
 - fever
 - flu-like symptoms (chills, body aches, feeling shaky, joint pain)
 - nausea and/or vomiting
 - dizziness or lightheadedness
 - fast or slow heart rate, or feeling like your chest is pounding
 - difficulty breathing or shortness of breath
 - changes in blood pressure

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection

How to store LEQEMBI IQLIK

Things to keep in mind when storing LEQEMBI IQLIK or taking it on the go



LEQEMBI IQLIK is safe at room temperature (77 °F/25 °C) for up to 14 days in the carton, but should not be placed back into the refrigerator once it reaches room temperature.



If you take your autoinjector out of the refrigerator, you can store it in a well-insulated container to ensure it doesn't get too hot or too cold. As long as it stays between 36 °F to 46 °F (2 °C to 8 °C), LEQEMBI IQLIK can go back into the refrigerator. **Do not try to speed up the warming process in any way.**

Do not take LEQEMBI IQLIK out of the carton or remove the cap until you're ready to inject.



Actor portrayal.
Actual LEQEMBI IQLIK carton shown.

Storage and handling of LEQEMBI IQLIK

- Store the LEQEMBI IQLIK autoinjector in a refrigerator between 36 °F to 46 °F (2 °C to 8 °C)
- Do not freeze
- Keep the LEQEMBI IQLIK autoinjector in the carton until ready to use, to protect it from light
- Do not shake LEQEMBI IQLIK
- LEQEMBI IQLIK may be stored at room temperature up to 77 °F (25 °C) for up to 14 days in the original carton. Do not place it back in the refrigerator if it has been brought to room temperature
- Throw away (dispose of) LEQEMBI IQLIK if it is out of date or not stored the right way
- Keep your LEQEMBI IQLIK autoinjector and all medicines out of the reach of children

IMPORTANT SAFETY INFORMATION (CONTINUED)

Infusion-related reactions:

- If you have an infusion-related reaction, your healthcare provider may give you medicines before your next infusion to lower the chance of having a reaction

Injection-related reactions:

- Injection-related reactions may occur with LEQEMBI injection under the skin (subcutaneous injection with LEQEMBI IQLIK). Tell your healthcare provider if you notice any of these symptoms during or after an injection:
 - redness, swelling, heat, pain, itching, rash, bruising, and blood collection under the skin at the injection site
 - headache, fatigue, or fever may also be observed after an injection

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection

Where to inject

LEQEMBI IQLIK can be injected into the front of the thighs, stomach area (abdomen), or back of the upper arms with the help of a loved one

You have options for where to inject—whichever is most comfortable for you.

Do not inject within 1 inch of your last injection location. Do not inject into moles, scars, bruises, or tattoos, or into areas where the skin is red, hard, tender, or injured.



**Stomach area
(abdomen)**

Do not inject within 2 inches of your belly button

**Back of the
upper arms**

With the help of a care partner

**Front of
the thighs**

For step-by-step instructions
on how to inject, visit [LEQEMBI.com/Injection](https://www.leqembi.com/Injection)

IMPORTANT SAFETY INFORMATION (CONTINUED)

The most common side effects of LEQEMBI include infusion-related reactions, ARIA, and headaches.

These are not all the possible side effects of LEQEMBI. Call your doctor for more information and medical advice about side effects. You may report side effects to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection

What to keep in mind when injecting



How to prepare

Once you are prescribed LEQEMBI IQLIK, you will likely receive a **practice autoinjector kit** (without medication) from your specialty pharmacy or healthcare provider. The kit may be delivered by mail or given in person.

The kit is a helpful way to get familiar with injecting, but it is not needed to safely use LEQEMBI IQLIK.

If you do not receive your kit, contact the LEQEMBI Companion™ program (see [page 11](#) for more information).



How to inject

When injecting, refer to the Instructions for Use. You can find it in the LEQEMBI IQLIK box or at LEQEMBI.com/Injection.



How to dispose of autoinjectors

Your specialty pharmacy may provide you with a sharps disposal container.

You may also be able to use a household container that is made of heavy-duty plastic and can be closed with a tight-fitting, puncture-resistant lid.

To learn more about safe sharps disposal and ways of getting a container, go to fda.gov/safesharpsdisposal.



How to get injection support

If you have questions about injecting, reach out to your healthcare provider.

You can also find helpful resources, including a video about how to use LEQEMBI IQLIK, at LEQEMBI.com/IFU.

IMPORTANT SAFETY INFORMATION (CONTINUED)

Before receiving LEQEMBI, tell your healthcare provider about:

- **All your medical conditions**, including if you are pregnant, breastfeeding, or plan to become pregnant or breastfeed. It is not known if LEQEMBI could harm your unborn or breastfeeding baby
- **All the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take medicines to reduce blood clots from forming (antithrombotic medicines, including aspirin)

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection

Understanding potential side effects

As your healthcare provider may have mentioned when you first started LEQEMBI infusions, **ARIA (amyloid-related imaging abnormalities)** is a potential side effect of treatments like LEQEMBI. ARIA does not usually cause any symptoms, but serious symptoms can occur. ARIA can be fatal. It's important for you to know about ARIA and what to ask your healthcare provider.

Likelihood of ARIA during maintenance treatment

ARIA is more likely to occur earlier in your treatment experience.

Symptoms of ARIA

ARIA commonly shows up as temporary swelling in areas of the brain that usually goes away over time. Small spots of bleeding in or on the surface of the brain can occur. Less often, larger areas of bleeding in the brain can occur.

Most people with ARIA don't have any symptoms. However, some people may notice:

- headache
- vision changes
- seizures
- confusion that gets worse
- nausea
- difficulty speaking
- dizziness
- difficulty walking
- muscle weakness

➤ **LEQEMBI can cause serious side effects, including serious allergic reactions and infusion-related reactions.**

The most common side effects of LEQEMBI include:

- infusion-related reactions
- swelling in areas of the brain, with or without small spots of bleeding in or on the surface of the brain (ARIA)
- headache

These are not all the possible side effects of LEQEMBI. For more information, ask your healthcare provider or pharmacist.

Keep track of how you're feeling

Each person may react differently to LEQEMBI. Share with your healthcare provider how you're feeling throughout treatment.

Patient information is accurate as of August 2025.
People featured have been compensated for their time.



NINA actual LEQEMBI patient

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection

Frequently asked questions



Are LEQEMBI IQLIK injections just as effective as my initial infusions?

Yes. Ongoing treatment with either infusions or injections keeps LEQEMBI working in your body.



Does LEQEMBI IQLIK deliver the same medicine that I received with infusions?

LEQEMBI IQLIK autoinjectors contain the same medicine as the LEQEMBI infusions you've received.



What if I miss an injection?

If you miss a dose of LEQEMBI IQLIK, inject the missed dose as soon as possible up to 6 days after the missed dose. Continue (resume) your next injection on your regular dosing schedule. See [page 11](#) to learn how the LEQEMBI Companion™ app can help you stay on track with your injection schedule.



If I decide at-home injections aren't for me, can I go back to infusions?

If you have concerns about at-home injections, talk with your healthcare provider. They can help make sure you receive LEQEMBI in the way that works best for you.



What is a sharps disposal container and how do I get one?

A sharps container is used to safely dispose of needles and other sharp objects. More information is available at fda.gov/safesharpsdisposal. You can request your own sharps container from your specialty pharmacy, or you may buy one from local pharmacies and medical supply stores.



Can I travel with LEQEMBI IQLIK?

Yes, you can take LEQEMBI IQLIK with you on the go. Before you leave on a trip, always check the autoinjector to make sure it hasn't expired. You should also consider getting a small sharps container to bring with you.

Reach out to your specialty pharmacy in advance of your trip to let them know your schedule and make sure you have enough autoinjectors. They can help you arrange shipping to your destination, if necessary.

If you are flying, reach out to your airline to see if you can carry injection devices on the plane. At the airport, notify the TSA screener that you are carrying an autoinjector. It's also a good idea to bring a note from your healthcare provider with you in case you are asked for one.

See [page 6](#) for additional information on how to store LEQEMBI IQLIK.



What if I have questions or need help with LEQEMBI IQLIK?

It's a good idea to reach out to your healthcare provider with any questions you have about using LEQEMBI IQLIK. There are also resources available to help you learn how to inject and stay on track with treatment, including step-by-step instructions, at LEQEMBI.com/IFU.

To get answers to other frequently asked questions,
visit LEQEMBI.com/FAQs

IMPORTANT SAFETY INFORMATION (CONTINUED)

LEQEMBI (lecanemab-irmb) is available as:

- Intravenous infusion: 100 mg/mL
- Subcutaneous injection: 200 mg/mL

Please see Important Safety Information throughout. Click [underlined links](#) for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection



Navigate treatment with support by your side

The LEQEMBI Companion™ program can help you:



Get information about accessing treatment



Understand your insurance coverage and potential out-of-pocket costs



Identify financial support programs, including the LEQEMBI Copay Assistance Program, if you are eligible for them

Ready to get started? Talk to your healthcare provider about enrolling in the LEQEMBI Companion program today. If you're already enrolled and have questions, call 1-833-453-7362 (1-833-4-LEQEMBI). For more information about the LEQEMBI Companion program and additional educational resources, visit LEQEMBI.com/PatientSupport.

Make the LEQEMBI Companion™ app your treatment companion

The LEQEMBI Companion app is your digital partner designed to help support you at every step of your treatment. It was developed in partnership with Medisafe's Medication Management app.

With the app you can find helpful information on **how to inject**, a tool for **tracking where and when you inject**, and other resources all in one place.



Scan the QR code to download the app or visit LEQEMBI.com/CompanionAppSignUp to learn more about what the app has to offer

Download the app



These steps will direct you to the Medisafe app. From there, you will select LEQEMBI as your treatment.

- 1 Scan the QR code, then tap the **pop-up banner**
- 2 Tap "**Get**" or "**Install**" to download Medisafe
- 3 Follow the prompts and tap "**Get Started**"
- 4 If prompted, enter the **verification code A4MW**

You do not need to be enrolled in the LEQEMBI Companion program to download and use the app.

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING**.

 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection



Need injection support?

Whether you want to know how to inject or just get more comfortable with the injection process, visit [LEQEMBI.com/HowToInject](https://www.leqembi.com/HowToInject) for helpful guidance.



Actor portrayal.

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.



2026 Eisai Inc. and Biogen.
LEQE-US5210 March 2026
All trademarks are the property of their respective owners.
LEQEMBI® is a registered trademark of Eisai R&D Management Co., Ltd.
LEQEMBI Companion™ and LEQEMBI IQLIK™ are trademarks of
Eisai R&D Management Co., Ltd.



 **LEQEMBI® IQLIK™**
(lecanemab-irmb)
Injection