



For weekly
self-injections

WHAT TO EXPECT with LEQEMBI IQLIK®

(pronounced "i-click")

Your LEQEMBI IQLIK dose may include 1 or 2 injections each, depending on your prescription.

WHAT IS LEQEMBI?

LEQEMBI is a prescription medicine used to treat people with early Alzheimer's disease, which includes mild cognitive impairment (MCI) or mild dementia stage of disease.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about LEQEMBI?
LEQEMBI can cause serious side effects, including: ARIA (Amyloid-Related Imaging Abnormalities). ARIA is a side effect that does not usually cause any symptoms, but serious symptoms can occur. ARIA can be fatal.

- ARIA commonly shows up as temporary swelling in areas of the brain that usually goes away over time
- Small spots of bleeding in or on the surface of the brain can occur

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.

 **LEQEMBI IQLIK®**
(lecanemab-irmb)
Injection

BEFORE YOU BEGIN

Write down important contacts

Use the space below to keep your healthcare provider's and specialty pharmacy's contact information in one spot. A specialty pharmacy is the type of pharmacy that fills your LEQEMBI IQLIK prescription. They help coordinate delivery and can answer questions about your prescription.

➤ **Remember to save your specialty pharmacy's name and number in your phone in case they call you with delivery updates.**

Healthcare provider

Name

Phone

Specialty pharmacy

Name

Phone

Get your treatment wallet card



The wallet card lets healthcare professionals know that you or your loved one is taking an anti-amyloid treatment, in case of an emergency. You can print the card by visiting [LEQEMBI.com/wallet](https://leqembi.com/wallet) or **scan the QR code** with your phone to add the card directly to your digital wallet.

Download the LEQEMBI Companion™ app



Track injections, explore resources, set appointment reminders, and more with the **LEQEMBI Companion app**. Visit page 11 for instructions on how to download the app.

IMPORTANT SAFETY INFORMATION (continued)

- Less often, larger areas of bleeding in the brain can occur
- Most people with ARIA don't have any symptoms. However, some people may notice: headache, confusion, dizziness, vision changes, nausea, difficulty walking, seizures, difficulty speaking, or muscle weakness

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.

GETTING STARTED with LEQEMBI IQLIK

This booklet is designed to support you as you begin or transition to treatment with LEQEMBI IQLIK. Inside, you'll find helpful information and tools to guide you through what to expect.

Find the treatment information that applies to you

LEQEMBI IQLIK comes in 2 different doses, depending on how and when you start treatment. To use this booklet, check off the box that matches your treatment path below. Then look for that icon on the tabs that follow to find information that applies to you. Content with no icon next to it applies to everyone starting with LEQEMBI IQLIK.

💬 **If you don't know your dose, ask your healthcare provider.**



Two 250-mg autoinjectors

- ☐ **I'M STARTING treatment with LEQEMBI IQLIK from the first day of treatment.** My dose is two 250-mg autoinjectors, once a week
- ☐ **I'M SWITCHING from intravenous (IV) infusions to LEQEMBI IQLIK injections during my first 18 months of treatment.** My dose is two 250-mg autoinjectors, once a week



One 360-mg autoinjector

- ☐ **I'M CONTINUING treatment with LEQEMBI IQLIK after 18 months for maintenance.** My dose is one 360-mg autoinjector, once a week

Your healthcare provider will guide you through the injection of LEQEMBI IQLIK for at least your first 2 weekly doses and answer any questions. After that, they will decide if it is appropriate for you or a care partner to inject LEQEMBI IQLIK.

- Some people have a gene called ApoE4 that may increase the risk of ARIA. Talk to your healthcare provider about testing to see if you have this gene

 **LEQEMBI IQLIK®**
(lecanemab-irmb)
Injection





“
Alzheimer’s
may test me,
**BUT IT WON’T
DEFINE ME.**”

LEONARD
LEQEMBI PATIENT SINCE 2023

ABOUT LEQEMBI IQLIK

LEQEMBI IQLIK is an autoinjector that you or a care partner uses to inject LEQEMBI under your skin. This is also called a subcutaneous injection. LEQEMBI IQLIK contains the same medicine as the LEQEMBI infusion, with comparable results and safety.

There are 2 different doses of LEQEMBI IQLIK
Depending on your dose, your autoinjector may look different.



Using LEQEMBI IQLIK

The steps on the following pages will go over how to properly store, inject, and dispose of LEQEMBI IQLIK. You can find the full Instructions for Use in the back pocket of this guide, and a video about using LEQEMBI IQLIK at LEQEMBI.com/IFUVideo.



Scan the QR code or visit LEQEMBI.com/IFUVideo for a video on how to use LEQEMBI IQLIK

Call your healthcare provider or go to the nearest hospital emergency room right away if you have any of the symptoms listed on page 2.

Serious allergic reactions:

Do not receive LEQEMBI if you have serious allergic reactions to LEQEMBI, LEQEMBI IQLIK, or any of the ingredients.

- Tell your healthcare provider and get medical help right away if you notice any symptoms during or after treatment with LEQEMBI, including: swelling of the face, lips, mouth, or tongue; itchy bumps on the skin, also known as hives; or difficulty breathing



IMPORTANT SAFETY INFORMATION (continued)

- You may be at a higher risk of developing bleeding in the brain if you take medicines to reduce blood clots from forming (antithrombotic medicines) while receiving LEQEMBI. Talk to your healthcare provider to see if any of the medicines you’re taking increase this risk
- Your healthcare provider will check for ARIA with MRI (magnetic resonance imaging) scans before you start LEQEMBI and during treatment
- You should carry information that says you are receiving LEQEMBI, which can cause ARIA, and that ARIA symptoms can look like stroke symptoms

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.

WHAT TO KNOW BEFORE INJECTING

Your LEQEMBI IQLIK dose may include 1 or 2 injections each, depending on your prescription. Your healthcare provider will guide you through the injection of LEQEMBI IQLIK for at least your first 2 weekly doses and answer any questions. After that, they will decide if it is appropriate for you or a care partner to inject LEQEMBI IQLIK.

Please bring your LEQEMBI IQLIK medication to your appointments unless other arrangements have been made. Bring your care partner if they will be helping with injections.



Practice injecting with your demo kit

Once your healthcare provider prescribes LEQEMBI IQLIK, you will likely receive a kit that contains an autoinjector without medication so you can practice injecting.

The kit may be included with this booklet, mailed to you by your specialty pharmacy, or given to you by your healthcare provider. The kit is a helpful way to get familiar with injecting, but it is not needed to safely use LEQEMBI IQLIK.

Receiving your LEQEMBI IQLIK

Your specialty pharmacy will most likely reach out to you with a phone call to verify your insurance coverage. They may also try to confirm where and when your medicine should be delivered.



When your specialty pharmacy first calls, it may be from an unknown number. Be sure to save the number so you can recognize future calls and texts.



Once your LEQEMBI IQLIK shipment is delivered, follow the instructions on the following page and Instructions for Use to store it properly.

IMPORTANT SAFETY INFORMATION (continued)

Infusion-related reactions:

- Infusion-related reactions are a common side effect that can occur during or after completion with LEQEMBI injection into a vein (intravenously), which can be serious. Tell your healthcare provider right away if you notice any of these symptoms: fever; flu-like symptoms (chills, body aches, feeling shaky, joint pain); nausea and/or vomiting; dizziness or lightheadedness; fast or slow heart rate, or feeling like your chest is pounding; difficulty breathing or shortness of breath; or changes in blood pressure
- If you have an infusion-related reaction, your healthcare provider may give you medicines before your next infusion to lower the chance of having a reaction

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.

HOW TO STORE LEQEMBI IQLIK

Your LEQEMBI IQLIK autoinjectors arrive in refrigerated cartons

Each carton will contain your full dose, which may be either 1 or 2 autoinjectors.



You should continue storing your cartons in a refrigerator between 36 °F to 46 °F (2 °C to 8 °C). Do not freeze. Keep in the original carton until ready to use to protect it from light. When you are ready to inject, take 1 carton out of the refrigerator and let it sit at room temperature for 20 minutes. Do not shake. **Do not try to speed up the warming process in any way.**



Whether you're traveling or bringing your medicine to your healthcare provider's office, it's important to remember that your LEQEMBI IQLIK carton can also be stored at room temperature (77 °F/25 °C) for up to 14 days. Do not place back into the refrigerator once it reaches room temperature.

Throw away LEQEMBI IQLIK that is out of date, or if it is not stored the right way. Keep your LEQEMBI IQLIK autoinjector and all medicines out of the reach of children.



Do not take LEQEMBI IQLIK out of the carton or remove the cap until you're ready to inject.

Injection-related reactions:

- Injection-related reactions may occur with LEQEMBI injection under the skin (subcutaneous injection with LEQEMBI IQLIK). Tell your healthcare provider if you notice any of these symptoms during or after an injection: redness, swelling, heat, pain, itching, rash, bump, bruising, and blood collection under the skin at the injection site; headache, chills, fatigue, or fever may also be observed after an injection

LEQEMBI IQLIK®
(lecanemab-irmb)
Injection



Tracking treatment
with LEQEMBI IQLIK



Tracking maintenance
treatment after 18 months

HOW TO INJECT LEQEMBI IQLIK

When injecting, refer to the Instructions for Use (IFU) found in your carton or in the pocket of this booklet. The Instructions for Use is a step-by-step guide to using LEQEMBI IQLIK.

Your healthcare provider will guide you through the injection of LEQEMBI IQLIK for at least your first 2 weekly doses and answer any questions.



You can also find a video of how to use LEQEMBI IQLIK by scanning the QR code

How to dispose of LEQEMBI IQLIK autoinjectors

Your specialty pharmacy may provide you with an FDA-approved sharps disposal container. You may also use a household container that is made of heavy-duty plastic and can be closed with a tight-fitting, puncture-resistant lid that's upright, leak-resistant, and properly labeled to warn of hazardous waste.




To learn more about safe sharps disposal and ways to get a container, go to [FDA.gov/SafeSharpsDisposal](https://www.fda.gov/SafeSharpsDisposal).

Where to inject

You have options for where to inject—whichever is most comfortable for you.

LEQEMBI IQLIK can be injected into the front of the thighs, stomach area (abdomen), or back of the upper arms with the help of a healthcare provider or a care partner.

Do not inject within 1 inch of last week's injection site. Do not inject into moles, scars, bruises, or tattoos, or into areas where the skin is red, hard, tender, or injured.



If your dose includes 2 autoinjectors, use the second autoinjector immediately after the first to complete your full prescribed dose. Use a different injection site at least 1 inch away from the last site each time you use LEQEMBI IQLIK.

IMPORTANT SAFETY INFORMATION (continued)

The most common side effects of LEQEMBI include infusion-related reactions with LEQEMBI injected into a vein, ARIA, and headache.

These are not all the possible side effects of LEQEMBI. Call your doctor for more information and medical advice about side effects. You may report side effects to the FDA at www.fda.gov/medwatch or call 1-800-FDA-1088.

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.



» For care partners

As a care partner, you can help inject LEQEMBI IQLIK in any of the highlighted locations. If your loved one would like to inject in the back of the arm, they will need your help.

You or a care partner can use the tracking pages toward the end of this booklet to track where and when you inject.

Before receiving LEQEMBI, tell your healthcare provider about:

- All your medical conditions, including if you are pregnant, breastfeeding, or plan to become pregnant or breastfeed. It is not known if LEQEMBI could harm your unborn or breastfeeding baby



Tracking treatment with LEQEMBI IQLIK



Tracking maintenance treatment after 18 months



Navigate treatment with support by your side

The LEQEMBI Companion™ program can help you:



Get information about accessing treatment



Understand your insurance coverage and potential out-of-pocket costs



Identify financial support programs, including the LEQEMBI Copay Assistance Program, if you are eligible for them

Ready to get started?

Talk to your healthcare provider about enrolling in the LEQEMBI Companion program today. If you're already enrolled and have questions, call **1-833-453-7362 (1-833-4-LEQEMBI)**.


For more information about the LEQEMBI Companion program and additional educational resources, visit LEQEMBI.com/PatientSupport

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.

Stay on track with the LEQEMBI Companion™ app

The LEQEMBI Companion app is your digital partner designed to help support you at every step of your treatment.

Use the app to start treatment and stay on track. It makes it easy to log your injections, set reminders, stay on top of your schedule, and more. Talk to your healthcare provider if you have any questions about your treatment plan.



Download the app



These steps will direct you to the Medisafe app. From there, you will select LEQEMBI as your treatment.

- 1 Scan the QR code, then tap the **pop-up banner**
- 2 Tap **"Get"** or **"Install"** to download Medisafe
- 3 Follow the prompts and tap **"Get Started"**
- 4 If prompted, enter the **verification code A4MW**



Scan the QR code to download the app or visit LEQEMBI.com/CompanionAppSignUp to learn more about what the app has to offer



Tracking treatment with LEQEMBI IQLIK



Tracking maintenance treatment after 18 months

MONITORING for potential side effects

It's important to know that all anti-amyloid treatments for Alzheimer's disease include MRI monitoring. LEQEMBI can cause serious side effects, including one called ARIA (Amyloid-Related Imaging Abnormalities). ARIA commonly shows up as temporary swelling or bleeding in the brain, and can only be seen with MRI monitoring.

Most of the time, ARIA does not cause any symptoms, though serious symptoms can occur and can be fatal.

That's why you will receive an MRI before treatment and 4 MRIs after you start treatment—to help your healthcare provider spot any changes early.

Your exact MRI schedule will be determined by your healthcare provider and may include additional MRIs.

Symptoms of ARIA

Most people with ARIA don't have any symptoms. However, some people may notice:

- headache
- vision changes
- seizures
- confusion
- nausea
- difficulty speaking
- dizziness
- difficulty walking
- muscle weakness

Likelihood of ARIA during treatment with LEQEMBI IQLIK



ARIA is more likely to occur earlier in your treatment experience. Your healthcare provider should schedule MRI exams after months 1, 2, 3, and 6 of treatment.



If you are switching to LEQEMBI IQLIK for maintenance injections, you will not need MRIs unless your healthcare provider requests them.

If ARIA does appear, your healthcare provider may pause or stop treatment and keep a closer eye on you. In most cases, ARIA resolves on its own.

LEQEMBI can cause serious side effects

These include serious allergic reactions and infusion- or injection-related reactions.

The most common side effects of LEQEMBI

These include infusion-related reactions with LEQEMBI injected into a vein, ARIA, and headache.

These are not all the possible side effects of LEQEMBI and LEQEMBI IQLIK. For more information, ask your healthcare provider or pharmacist.

Please see Important Safety Information throughout. Click underlined links for **Medication Guide with Instructions for Use** and **full Prescribing Information**, including **Boxed WARNING**.

TRACKING TREATMENT with LEQEMBI IQLIK



2 autoinjectors
per dose

Once
weekly

The following pages are designed to help you track your **treatment dates** and **injection locations** if you are:

- **starting with LEQEMBI IQLIK from the first day of treatment.** Your dose is two 250-mg autoinjectors, once a week
- **switching from IV infusions to LEQEMBI IQLIK injections during your first 18 months of treatment.** Your dose is two 250-mg autoinjectors, once a week



Prefer to track treatment on your phone?

Visit LEQEMBI.com/LEQEMBICompanion to learn how to download the LEQEMBI Companion™ app

LEQEMBI IQLIK®
(lecanemab-irmb)
Injection



Tracking treatment
with LEQEMBI IQLIK




Tracking maintenance
treatment after 18 months

REMEMBER: 2 LEQEMBI IQLIK autoinjectors is 1 complete dose

STAYING ON TRACK with TREATMENT

Your MRI schedule at a glance

The 4 MRIs your healthcare provider schedules after starting treatment with LEQEMBI IQLIK will help monitor your brain and check for changes early. They will take place after months 1, 2, 3 and 6.



Your exact MRI schedule will be determined by your healthcare provider and may include additional MRIs.

As you're going through treatment, MRIs help your healthcare provider monitor how your body is responding and ensure it's safe to continue.

Some people taking LEQEMBI experience a side effect called ARIA, which commonly shows up as temporary swelling or bleeding in the brain that can be fatal. Most of the time, ARIA does not cause any symptoms, and people don't feel any different—it may show up as small spots that can only be seen with an MRI. In most cases, ARIA usually resolves on its own.



Ask your healthcare provider if you can schedule MRIs ahead of time.

Your healthcare provider may follow up with you to discuss your MRI results.

If you are switching from IV infusions to self-injections, you may have already completed some MRIs.

Talk to your healthcare provider to see where you are in the MRI schedule.

IMPORTANT SAFETY INFORMATION (continued)

Before receiving LEQEMBI, tell your healthcare provider about:

- **All the medicines you take**, including prescription and over-the-counter medicines, vitamins, and herbal supplements. Especially tell your healthcare provider if you take medicines to reduce blood clots from forming (antithrombotic medicines, including aspirin)

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including Boxed WARNING.

REMEMBER: 2 LEQEMBI IQLIK autoinjectors is 1 complete dose



MY LEQEMBI IQLIK SELF-INJECTION LOG

Use the tracker on the following pages to log the date, time, and location of your once-weekly, 2-autoinjector dose.

TIP: Picking a specific day of the week to do your self-injection may help with staying on track.

It's important to stick to your treatment schedule.

Your injections or infusions keep LEQEMBI working in your body and fighting Alzheimer's disease.

If you miss a scheduled injection, inject both autoinjectors as soon as you are able. You can still inject up to 6 days after the missed dose. Then take your next dose on your usual scheduled day. After that, you can continue with your normal dosing schedule. Visit [LEQEMBI.com/HowToInject](https://www.leqembi.com/HowToInject) for an overview of the injection process.

LEQEMBI (lecanemab-irmb) is available as:

- Intravenous infusion: 100 mg/mL
- Subcutaneous injection: 200 mg/mL

LEQEMBI IQLIK®
(lecanemab-irmb)
Injection



REMEMBER: 2 LEQEMBI IQLIK autoinjectors is 1 complete dose

MY SELF-INJECTIONS

Where to inject

Back of the upper arms
With the help of a care partner

Stomach area (abdomen)
Do not inject within 2 inches of your belly button

Front of the thighs

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time



If your dose includes 2 autoinjectors, make sure your injections are at least 1 inch apart. Use the second autoinjector immediately after the first to complete your full prescribed dose.

REMEMBER: 2 LEQEMBI IQLIK autoinjectors is 1 complete dose

MY SELF-INJECTIONS

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time

Week #

Date

Time




If you choose to inject into the back of the upper arms, you will need the help of a healthcare provider or care partner.

LEQEMBI IQLIK®
(lecanemab-irmb)
Injection



REMEMBER: 2 LEQEMBI IQLIK autoinjectors is 1 complete dose

MY SELF-INJECTIONS


	Week # Date Time
	Week # Date Time
	Week # Date Time

If you've finished 18 months of treatment with LEQEMBI IQLIK and talked with your healthcare provider about continuing treatment, you can use the next section of this booklet to track your maintenance doses.



If you choose to inject into the back of the upper arms, you will need the help of a healthcare provider or care partner.

**Download more
self-injection log pages**
by visiting
LEQEMBI.com/tracker



Prefer to track treatment on your phone?
Scan the QR code to download the app or
visit LEQEMBI.com/CompanionAppSignUp

 **LEQEMBI IQLIK®**
(lecanemab-irmb)
Injection



Tracking maintenance
treatment after 18 months

NOTES



TRACKING MAINTENANCE TREATMENT with LEQEMBI IQLIK



1 autoinjector
per dose

Once
weekly

The following pages are designed to help you track your **treatment dates** and **injection locations** if you are:

- **continuing treatment with LEQEMBI IQLIK for maintenance.** Your dose is one 360-mg autoinjector, once a week

» Prefer to track treatment on your phone?

Visit LEQEMBI.com/LEQEMBICompanion to learn how to download the LEQEMBI Companion™ app

LEQEMBI IQLIK®
(lecanemab-irmb)
Injection



REMEMBER: 1 LEQEMBI IQLIK autoinjector is 1 complete dose

WHAT IS MAINTENANCE TREATMENT?

After treating with LEQEMBI for 18 months, you may have the option of starting maintenance treatment. However, you and your healthcare provider may decide to continue your current treatment plan, even after 18 months. Talk to your healthcare provider to determine the plan that's best for you.

Only LEQEMBI has FDA-approved maintenance dosing options to keep slowing progression.



If you are beginning maintenance treatment with LEQEMBI IQLIK, your dose is one 360-mg autoinjector, once weekly (every 7 days).

If this is your first time using LEQEMBI IQLIK, your healthcare provider will guide you through the injection of LEQEMBI IQLIK for at least your first 2 weekly doses and answer any questions.

It's important to stick to your treatment schedule.

Your injections keep LEQEMBI working in your body and fighting Alzheimer's disease.

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about LEQEMBI? LEQEMBI can cause serious side effects, including: ARIA (Amyloid-Related Imaging Abnormalities). ARIA is a side effect that does not usually cause any symptoms, but serious symptoms can occur. ARIA can be fatal.

- ARIA commonly shows up as temporary swelling in areas of the brain that usually goes away over time

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#)

and [full Prescribing Information](#), including Boxed WARNING.

REMEMBER: 1 LEQEMBI IQLIK autoinjector is 1 complete dose



MY LEQEMBI IQLIK SELF-INJECTION LOG

Use the tracker on the following pages to log the date, time, and location of your once-weekly, 1-autoinjector dose.

Back of the upper arms
With the help of a care partner

Date TUES 7/15

Time 2:30pm

This should be 1 week from your previous injection

Circle the location on your body where you used the autoinjector

TIP: Picking a specific day of the week to do your self-injection may help with staying on track.

It's important to stick to your treatment schedule.

Your injections keep LEQEMBI working in your body and fighting Alzheimer's disease.

If you miss a scheduled injection, inject as soon as you remember. You can still inject up to 6 days after the missed dose. Then take your next dose on your usual scheduled day. After that, you can continue with your normal dosing schedule. Visit [LEQEMBI.com/HowToInject](https://www.leqembi.com/HowToInject) for an overview of the injection process.


- Small spots of bleeding in or on the surface of the brain can occur
- Less often, larger areas of bleeding in the brain can occur
- Most people with ARIA don't have any symptoms. However, some people may notice: headache, confusion, dizziness, vision changes, nausea, difficulty walking, seizures, difficulty speaking, or muscle weakness

 **LEQEMBI IQLIK®**
(lecanemab-irmb)
Injection

REMEMBER: 1 LEQEMBI IQLIK autoinjector is 1 complete dose

MY SELF-INJECTIONS

Where to inject




Back of the upper arms
With the help of a care partner

Stomach area (abdomen)
Do not inject within 2 inches
of your belly button

Front of the thighs

Download more
self-injection log pages
by visiting
LEQEMBI.com/tracker



Prefer to track treatment on your phone?
Scan the QR code to download the app or
visit LEQEMBI.com/CompanionAppSignUp

 **LEQEMBI IQLIK®**
(lecanemab-irmb)
Injection



If you choose to inject into the back of the upper arms, you will need the help of a healthcare provider or care partner.

NOTES

What to expect with
LEQEMBI IQLIK

Tracking treatment
with LEQEMBI IQLIK



Tracking maintenance
treatment after 18 months



LANCE

LEQEMBI IQLIK PATIENT

“
I SEE A FUTURE
*that will continue to make
important progress in
the treatment of
Alzheimer’s disease.*

People shown were compensated for their time, and information is accurate as of February 2026. Quotes reflect personal experiences, which may vary.



Get tips for care partners
by visiting LEQEMBI.com/CarePartners

 **LEQEMBI IQLIK[®]**
(lecanemab-irmb)
Injection

“

I'm living proof that
**ACTING EARLY CAN
MAKE A DIFFERENCE.**

LEONARD

LEQEMBI IQLIK PATIENT

People shown were compensated for their time, and information is accurate as of February 2026. Quotes reflect personal experiences, which may vary.



Scan the QR code or visit LEQEMBI.com/IFUVideo
for step-by-step instructions on how to inject

IMPORTANT SAFETY INFORMATION (continued)

What is the most important information I should know about LEQEMBI?

- Some people have a gene called ApoE4 that may increase the risk of ARIA. Talk to your healthcare provider about testing to see if you have this gene
- You may be at a higher risk of developing bleeding in the brain if you take medicines to reduce blood clots from forming (antithrombotic medicines) while receiving LEQEMBI. Talk to your healthcare provider to see if any of the medicines you're taking increase this risk

Please see Important Safety Information throughout. Click underlined links for [Medication Guide with Instructions for Use](#) and [full Prescribing Information](#), including **Boxed WARNING.**

 **LEQEMBI IQLIK®**
(lecanemab-irmb)
Injection



© 2026 Eisai Inc. and Biogen.

LEQE-US5290 July 2026

All trademarks are the property of their respective owners.

LEQEMBI® and LEQEMBI IQLIK® are registered trademarks of

Eisai R&D Management Co., Ltd.

LEQEMBI Companion™ is a trademark of Eisai R&D Management Co., Ltd.